

12 Radiation from Relativistic Charged Particles

12.1 Basic equations

(a) We wrote down the wave equations in the covariant gauge:

$$-\square\varphi = \rho(t_o, \mathbf{r}_o) \quad (12.1)$$

$$-\square\mathbf{A} = \mathbf{J}(t_o, \mathbf{r}_o)/c \quad (12.2)$$

(b) Then we used the green function of the wave equation

$$G(t, r|t_o r_o) = \frac{1}{4\pi|\mathbf{r} - \mathbf{r}_o|} \delta(t - t_o + \frac{|\mathbf{r} - \mathbf{r}_o|}{c}) \quad (12.3)$$

to determine the potentials (φ, \mathbf{A}) with the current

$$\frac{J^\mu}{c} = (\rho, \frac{\mathbf{J}}{c}) = (q\delta^3(\mathbf{r}_o - \mathbf{r}_*(t_o)), q\frac{\mathbf{v}(t_o)}{c}\delta^3(\mathbf{r}_o - \mathbf{r}_*(t_o))) \quad (12.4)$$

This yields the Lienard-Wiechert potentials

$$\varphi = \frac{q}{4\pi|\mathbf{r} - \mathbf{r}_*(T)|} \frac{1}{1 - \mathbf{n} \cdot \boldsymbol{\beta}(T)} \implies \frac{q}{4\pi r} \frac{1}{1 - \mathbf{n} \cdot \boldsymbol{\beta}(T)} \quad (12.5)$$

$$\mathbf{A} = \frac{q}{4\pi|\mathbf{r} - \mathbf{r}_*(T)|} \frac{\mathbf{v}(T)/c}{1 - \mathbf{n} \cdot \boldsymbol{\beta}(T)} \implies \frac{q}{4\pi r} \frac{\mathbf{v}(T)/c}{1 - \mathbf{n} \cdot \boldsymbol{\beta}(T)} \quad (12.6)$$

where the retarded time is

$$T(t, r) = t - \frac{|\mathbf{r} - \mathbf{r}_*(T)|}{c} \implies T(t, r) = t - \frac{r}{c} + \frac{\mathbf{n} \cdot \mathbf{r}_*(T)}{c} \quad (12.7)$$

The terms after the Longrightarrow indicate the far field limit

(c) The Lienard Wiechert potential can also be obtained by integrating over \mathbf{r}_o in Eq. (10.8).

(d) The factor ‘‘collinear factor’’ (my name), or dT/dt

$$\frac{dT}{dt} = \frac{1}{(1 - \mathbf{n} \cdot \boldsymbol{\beta})} \quad (12.8)$$

$$\frac{dT}{dr^i} = \frac{1}{(1 - \mathbf{n} \cdot \boldsymbol{\beta})} \frac{-n_i}{c} \quad (12.9)$$

is quite important. We gave a physical interpretation of it in class. If a wave form is *observed* to have a time scale of Δt , then the *formation time* of the wave, ΔT , is

$$\Delta T = \frac{dT}{dt} \Delta t = \frac{\Delta t}{1 - \mathbf{n} \cdot \boldsymbol{\beta}} \quad (12.10)$$

In particular, a fourier component with frequency ω in the observed wave was formed over the time

$$\Delta T \sim \frac{1}{\omega(1 - \mathbf{n} \cdot \boldsymbol{\beta})} \quad (12.11)$$

- (e) The magnetic and electric fields can be determined from $\mathbf{E} = -\frac{1}{c}\partial_t\mathbf{A}_{\text{rad}} - \nabla\varphi$. As discussed in a separate note (“retarded_time.pdf”), In the far field limit this is the same as computing

$$\mathbf{E}(t, r) = \mathbf{n} \times \mathbf{n} \times \frac{1}{c}\partial_t\mathbf{A}_{\text{rad}}(T) \quad (12.12a)$$

$$= \mathbf{n} \times \mathbf{n} \times \frac{1}{1 - \mathbf{n} \cdot \boldsymbol{\beta}} \frac{1}{c} \frac{\partial}{\partial T} \mathbf{A}_{\text{rad}}(T) \quad (12.12b)$$

$$= \frac{1}{1 - \mathbf{n} \cdot \boldsymbol{\beta}} \frac{\partial}{\partial T} \left[\frac{q}{4\pi r c^2} \frac{\mathbf{n} \times \mathbf{n} \times \mathbf{v}/c}{1 - \mathbf{n} \cdot \boldsymbol{\beta}} \right]_{\text{ret}} \quad (12.12c)$$

$$= \frac{q}{4\pi r c^2} \left[\frac{\mathbf{n} \times (\mathbf{n} - \boldsymbol{\beta}) \times \mathbf{a}}{(1 - \mathbf{n} \cdot \boldsymbol{\beta})^3} \right]_{\text{ret}} \quad (12.12d)$$

The $[\]_{\text{ret}}$ indicates that the velocity and acceleration are to be evaluated at the retarded time $T(t, r)$.

The magnetic field is

$$\mathbf{B} = \mathbf{n} \times \mathbf{E} \quad (12.13)$$

- (f) We will often be interested in the frequency distribution of the radiation. Computing the fourier transform of \mathbf{E} yields straightforwardly with Eq. (12.12) and the collinear factor, Eq. (12.8)

$$\mathbf{E}(\omega, r) = \int_{-\infty}^{\infty} e^{i\omega t} \mathbf{E}(t, r) \quad (12.14)$$

$$= \frac{q(-i\omega e^{i\omega r/c})}{4\pi r c^2} \int_{-\infty}^{\infty} dT e^{i\omega(T - \mathbf{n} \cdot \mathbf{r}_*(T)/c)} \mathbf{n} \times \mathbf{n} \times \mathbf{v}(T)/c \quad (12.15)$$

This final form is often the most convenient, but sometimes it is just easier to use

$$\mathbf{E}(\omega, r) = \frac{q e^{i\omega r/c}}{4\pi r c^2} \int_{-\infty}^{\infty} dT e^{i\omega(T - \mathbf{n} \cdot \mathbf{r}_*(T)/c)} \frac{\mathbf{n} \times (\mathbf{n} - \boldsymbol{\beta}) \times \mathbf{a}}{(1 - \mathbf{n} \cdot \boldsymbol{\beta})^2} \quad (12.16)$$

which shows explicitly the dependence on acceleration

Observables in the far field

- (a) The energy per time per solid angle received at the detector is

$$\frac{dW}{dt d\Omega} = \frac{dP(t)}{d\Omega} = r^2 \mathbf{S} \cdot \mathbf{n} \quad (12.17)$$

$$= c|r\mathbf{E}|^2 \quad (12.18)$$

This is what you want to know if you want to find out if the detector will burn up.

- (b) We often want to know how much energy was radiated over a given period of acceleration, $T_1 \dots T_2$. For example how much energy was lost by the particle as it moved through one complete circle. Then we want to evaluate the energy radiated per retarded time from T_1 up to the time it completes the circle T_2

$$\frac{dW}{dT d\Omega} = \frac{dP(T)}{d\Omega} = r^2 \mathbf{S} \cdot \mathbf{n} \frac{dt}{dT} \quad (12.19)$$

$$= c|r\mathbf{E}|^2 (1 - \mathbf{n} \cdot \boldsymbol{\beta}) \quad (12.20)$$

- (c) We are also interested in the frequency distribution of the emitted radiation. The energy per $d\omega/(2\pi)$ per solid angle is

$$(2\pi) \frac{dW}{d\omega d\Omega} \equiv c|rE(\omega, r)|^2 \quad (12.21)$$

Since the sign of the ω is without significance (for real fields such as the electromagnetic fields), we sometimes use

$$\frac{dI}{d\omega d\Omega} \equiv \frac{c|rE(\omega, r)|^2}{2\pi} + \frac{c|rE(-\omega, r)|^2}{2\pi} = \frac{c|rE(\omega, r)|^2}{\pi} \quad (12.22)$$

So that

$$\frac{dW}{d\Omega} = \int_0^\infty \frac{dI}{d\omega d\Omega} \quad (12.23)$$

- (d) The energy spectrum can be interpreted as the average number of photons per frequency per solid angle

$$\frac{dI}{d\omega d\Omega} = \hbar\omega \frac{d\bar{N}}{d\omega d\Omega} \quad (12.24)$$

12.2 Relativistic Larmour

- (a) For a particle undergoing arbitrary relativistic motion, we evaluated the energy per retarded time per solid angle

$$\frac{dP(T)}{d\Omega} = \frac{q^2}{16\pi^2 c^3} \frac{|\mathbf{n} \times (\mathbf{n} - \boldsymbol{\beta}) \times \mathbf{a}|^2}{(1 - \mathbf{n} \cdot \boldsymbol{\beta})^5} \quad (12.25)$$

- (b) Integrating over angles we get

$$P(T) = \frac{dW}{dT} = \frac{q^2}{4\pi} \frac{2}{3c^3} \gamma^6 \left[a_{\parallel}^2 + \frac{a_{\perp}^2}{\gamma^2} \right] \quad (12.26)$$

where a_{\parallel} is the projection of $\mathbf{a} = d^2\mathbf{x}/dt^2$ along the direction of motion, and a_{\perp} is the component of \mathbf{a} perpendicular to the direction of motion, *i.e.* for \mathbf{v} in the z direction

$$\mathbf{a} = (a_{\perp}^x, a_{\perp}^y, a_{\parallel}) \quad (12.27)$$

- (c) The acceleration four vector is

$$\mathcal{A}^{\mu} = \frac{d^2 x^{\mu}}{d\tau^2} \quad (12.28)$$

For a particle moving along in the z -direction, the acceleration in the particle's locally inertial frame (*i.e.* the frame that is instantaneously moving with the particle) is

$$(\mathcal{A}^0, \mathcal{A}^1, \mathcal{A}^2, \mathcal{A}^3)|_{\text{rest frame}} = (0, \alpha_{\perp}^x, \alpha_{\perp}^y, \alpha_{\parallel}) \quad (12.29)$$

While in the lab frame \mathcal{A}^{μ} is found by boosting this result. The acceleration $\mathbf{a} = \frac{d\mathbf{v}}{dt}$ is found from this result and the definition of proper time $d\tau = dt/\gamma$,

$$\mathbf{a} = (a_{\perp}^x, a_{\perp}^y, a_{\parallel}) = (\gamma^2 \alpha_{\perp}^x, \gamma^2 \alpha_{\perp}^y, \gamma^3 \alpha_{\parallel}) \quad (12.30)$$

You should be able to prove this. The relativistic Larmour formula can then be written

$$P(T) = \frac{q^2}{4\pi} \frac{2}{3c^3} \mathcal{A}_{\mu} \mathcal{A}^{\mu} \quad (12.31)$$

- (d) For straight line acceleration at very large γ , we found that that the radiation is emitted within a cone of order

$$\Delta\Theta \sim 1/\gamma. \quad (12.32)$$

For θ very small $\theta \sim 1/\gamma$ we found,

$$\frac{dP(T)}{d\Omega} = \frac{2q^2}{\pi^2} \frac{a^2}{c^3} \gamma^8 \frac{(\gamma\theta)^2}{(1 + (\gamma\theta)^2)^5}. \quad (12.33)$$

You should feel comfortable deriving this result.