


# Legendre Polynomials

For azimuthally symmetric problems don't need all  $Y_{lm}$ . Since  $Y_{lm} \propto e^{im\phi}$  need only  $m=0$ .


$$Y_{l0} = \sqrt{\frac{2l+1}{4\pi}} P_l(\cos\theta)$$

a polynomial in  $\cos\theta$

Any function of  $\theta$  can be expanded in Legendre polynomials (see handout):

$$(1) \quad f(\cos\theta) = \sum_l f_l \left(\frac{2l+1}{2}\right) P_l(\cos\theta) \quad (\text{expansion})$$

(2) Orthogonal

$$\int_{-1}^1 d(\cos\theta) P_l(\cos\theta) P_{l'}(\cos\theta) = \frac{2}{2l+1} \delta_{ll'}$$

$$(3) \quad f_l = \int_{-1}^1 d\cos\theta P_l(\cos\theta) f(\cos\theta) \quad \text{coefficient}$$

$$(4) \quad \int_{-1}^1 P_l(\cos\theta) P_{l'}(\cos\theta) \frac{2l+1}{2} = \delta(\cos\theta - \cos\theta')$$

$$P_0 = 1 \quad P_2 = \frac{1}{2}(3x^2 - 1)$$

$$P_1 = x$$


## Legendre Polynom

Similarly, for azimuthally symmetric systems

$$\phi(\vec{r}) = \sum_{l=0}^{\infty} (A_l r^l + \frac{B_l}{r^{l+1}}) P_l(\cos \theta)$$

## In class problem pg 1

- A neutral metal sphere of radius  $a$  is placed in an electric field



Determine the potential Everywhere

### Solution

- Boundary Conditions  $\varphi = \text{const}$  on surface, and  $\varphi \xrightarrow{r \rightarrow \infty} -E_0 z = -E_0 r \cos\theta + \varphi_0$    
  $\varphi_0$  could set to zero
- Outside:

$$\varphi(r) = \sum_l (A_l^{\text{out}} r^l + \frac{B_l^{\text{out}}}{r^{l+1}}) P_l(\cos\theta)$$

$r \rightarrow \infty$  Boundary conditions:  $A_l^{\text{out}} = 0$ , Except  $A_1^{\text{out}}$

$$A_1^{\text{out}} = -E_0 r \cos\theta, \text{ and in principle } A_0^{\text{out}} = \varphi_0$$

$$\varphi^{\text{out}} = -E_0 r \cos\theta + \sum_l \frac{B_l^{\text{out}}}{r^{l+1}} P_l(\cos\theta) + \varphi_0$$

# Sphere - In Class pg 2

Now since  $\varphi$  is continuous,

$$\varphi^{\text{out}} = \text{const} \quad \text{as } r \rightarrow a + \text{bit} \quad \text{tiny bit}$$

Says that  $B_\ell = 0$  unless  $\ell = 1$   $m = 0$

$$\varphi^{\text{out}} = \varphi_0 + -E_0 r \cos\theta + \frac{B_1}{r^2} \overbrace{P_1(\cos\theta)}$$

$$\varphi^{\text{out}} = \varphi_0 + -E_0 r \cos\theta + \frac{B}{r^2} \cos\theta$$

Requiring that  $\varphi = \text{const}$  at  $r = a$ , sets  $B = a^3 E_0$

$$\varphi^{\text{out}} = \varphi_0 - E_0 r \cos\theta + \frac{a^3 E_0}{r^2} \cos\theta$$

Similarly for  $r < a$

$$\varphi^{\text{in}} = \sum_{\ell m} \left( A_\ell r^\ell + \frac{B_{\ell m}}{r^{\ell+1}} \right) P_\ell$$

(regularity)

Then since  $\varphi|_{r=a} = \text{const}$ , and continuity gives

$$\varphi^{\text{in}} = \varphi_0$$

So

$$\varphi(r, \cos\theta) = \begin{cases} \varphi_0 - E_0 r \cos\theta + \frac{a^3 E_0 \cos\theta}{r^2} & \text{outside} \\ \varphi_0 & \text{inside} \end{cases}$$

The charge density:


$$E_r^{\text{out}} - E_r^{\text{in}} = \sigma$$

$$-\frac{\partial \varphi^{\text{out}}}{\partial r} + \frac{\partial \varphi^{\text{in}}}{\partial r} = \sigma$$

$$E_0 \cos\theta + \frac{2a^3 E_0 \cos\theta}{r^3} \Big|_{r=a} = \sigma$$

$$\boxed{3E_0 \cos\theta = \sigma}$$

So picture



To determine the dipole moment, compare the potential

$$\varphi_{\text{ind}} = \frac{1}{4\pi r} Q_{\text{tot}} + \frac{\vec{p} \cdot \hat{r}}{4\pi r^2} + \dots$$

To our potential.

$$\varphi(\vec{r}) = \underbrace{\varphi_0 - E_0 r \cos\theta}_{\text{external field}} + \underbrace{\frac{a^3 E_0 \cos\theta}{r^2}}_{\text{induced field}}$$

So,  $\vec{p} = 4\pi a^3 E_0 \hat{z}$  by comparison of

$$\frac{\vec{p} \cdot \hat{r}}{4\pi r^2} \quad \text{and} \quad \frac{a^3 E_0 \cos\theta}{r^2}$$