


Canonical Transformations

- We want to find new coordinates for phase space.

- Why?



We already discussed that the motion of a SHO is a circle. The motion of a perturbed oscillator is close to a circle. Maybe we should use circular coordinates (Amplitude and Phase)? Or perhaps $a_{\pm} = wq \pm ip$?

- Change of coordinates which preserve the form of Hamilton's Equations are particularly important.

- So we look for a map

$$\left. \begin{aligned} q &\rightarrow Q(q, p) \\ p &\rightarrow P(q, p) \\ h(q, p, t) &\rightarrow \underline{H}(Q, P, t) \end{aligned} \right\} \begin{aligned} &\text{Where the new} \\ &\text{Eom are:} \\ &\dot{Q} = \partial \underline{H} / \partial P \\ &\dot{P} = -\partial \underline{H} / \partial Q \end{aligned}$$

- Then the action in the old coordinates is

$$S_1 = \int p dq - h dt$$

The action in the new coordinates is

$$S_2 = \int P dQ - \underline{H} dt \quad \leftarrow \text{the EOM for } Q, P \text{ are unchanged}$$

- The difference in the two actions can only be a total derivative, which modifies the boundary terms without modifying the EOM.

$$S_1 - S_2 = \int_{t_1}^{t_2} \frac{dF}{dt} dt$$

or taking $F(q, Q, t)$ we find

$$\begin{aligned} \int p dq - P dQ - (h - \underline{H}) dt \\ = \int \underbrace{\frac{\partial F}{\partial q} dq + \frac{\partial F}{\partial Q} dQ + \frac{\partial F}{\partial t} dt}_{dF/dt} \end{aligned}$$

So if we compare dF/dt

- So we compare both sides

Yielding

$$(1) \quad p = \frac{\partial F}{\partial q}(q, Q, t)$$

$$\text{and} \quad \underline{H} = h + \frac{\partial F}{\partial t}$$

$$(2) \quad -P = \frac{\partial F}{\partial Q}(q, Q, t)$$

- Gives a prescription: Use (1) to find $Q(q, p, t)$. then we can evaluate $P(q, Q(q, p, t))$ from (2).

- Rather than working with F it is easier to Legendre transform / integrate by parts

$$\begin{aligned} S_1 - S_2 &= \int p dq - P dQ - (h - \underline{H}) dt \\ &= \int dt \frac{dF}{dt}(q, Q, t) \end{aligned}$$

- Write $-P dQ = -d(PQ) + Q dP$, and bring it to the other side:

$$\begin{aligned} S_1 - S_2 &= \int p dq + Q dP - (h - \underline{H}) dt \\ &= \int \frac{d}{dt} (F + PQ) \end{aligned}$$

Then \swarrow also called F_2

$\Phi \equiv F + PQ$ is the Legendre Transform of F

And generates the following canonical map

$$d\bar{\Phi}(q, P) = p dq + Q dP - (H - \underline{H}) dt$$

i.e.

$$(1) \quad p = \frac{\partial \bar{\Phi}(q, P)}{\partial q}$$

$$\underline{H}(Q, P) = h(q, p) + \frac{\partial \bar{\Phi}}{\partial t}$$

$$(2) \quad Q = \frac{\partial \bar{\Phi}(q, P)}{\partial P}$$


- Works like before first we solve for $P(q, p)$ from (1), then evaluate $Q(q, p)$ from (2)

Examples

- $F = qQ$ exchanges q and p

$$p = \frac{\partial F}{\partial q} = Q$$

$$-P = \frac{\partial F}{\partial Q} = q$$



- Take $\bar{\Phi} = qP$, ^{this is the} identity transformation

$$p = \frac{\partial \bar{\Phi}}{\partial q} = P$$

$$Q = \frac{\partial \bar{\Phi}}{\partial P} = q$$

- Now that we know the identity transformation we can use a transformation close to the identity

$$\Phi = qP + G(q, P) \lambda \quad \leftarrow \text{small}$$

Then

$$(1) \quad p = P + \frac{\partial G}{\partial q}(q, P) \lambda$$

$$(2) \quad Q = q + \frac{\partial G}{\partial P}(q, P) \lambda$$

- Solving (1) by iteration. $p \approx P$ at zeroth order. Then we can substitute $p = P$ in $\partial G(q, P) / \partial q$ at first order:

$$\star \quad P \approx p - \frac{\partial G}{\partial q}(q, p) \lambda$$

$$Q = q + \frac{\partial G}{\partial P}(q, P) \lambda$$

$$\star \quad Q \approx q + \frac{\partial G}{\partial P}(q, p) \lambda$$

These two transformations are the infinitesimal transforms discussed first


Circular Coordinates in Phase Space, The SHO.

- The Hamiltonian of the SHO

$$H = \frac{1}{2} p^2 + \omega^2 q^2$$

- Is it possible to find a canonical transform where one of my coordinates P is the Hamiltonian itself? say

$$H = \omega P$$



This is an (amplitude)² and the phase representation of the Harmonic oscillator. $P = (\text{amplitude})^2$ and $Q = \text{phase}$. In the coordinates the solution is simple!


$$\dot{Q} = \frac{\partial H}{\partial P} = \omega \Rightarrow Q = \omega t + t_0$$

$$\dot{P} = -\frac{\partial H}{\partial Q} = 0 \Rightarrow P = \text{constant}$$

A healthy dose of numerology / guess work gives

$$F(q, Q) = \frac{\omega q^2}{2} \tan Q$$

- Let's de-mystify this transformation a bit:



$$P = (\text{radius})^2 \propto \text{energy of SHO}$$

- Want:

$$\frac{P}{wq} = \tan Q = \tan(\text{phase})$$

$$P = \frac{1}{2} wq^2 / \cos^2 Q = \frac{1}{2} w \underbrace{(q / \cos Q)^2}_{(\text{amplitude})^2}$$

Then we want

$$(\star) \quad p = wq \tan Q = \frac{\partial F}{\partial q}$$

$$(\star\star) \quad P = \frac{1}{2} \frac{wq^2}{\cos^2 Q} = \frac{\partial F}{\partial Q}$$

- What is non-trivial about the transformation is it is "curl free":

$$\frac{\partial p}{\partial Q} = \frac{\partial P}{\partial q} \quad \text{or} \quad \frac{\partial^2 F}{\partial q \partial Q} = \frac{\partial^2 F}{\partial Q \partial q}$$

- Then this guarantees, that the desired transformation could be integrated to find the generating function.

$$F(q, Q) = wq^2 \tan Q / 2$$

So we can use circular coordinates for phase space

$$Q = \tan \frac{p}{\omega q}$$

$$P = \frac{1}{2} p^2 + \frac{1}{2} (\omega q)^2$$

The EOM in these coordinates will always be the same $\dot{Q} = \partial H / \partial P$ $\dot{P} = -\partial H / \partial Q$. Using this coordinate system is useful for problems which are close to the SHO