



A worked example pg. 1

Charged Cylinder: $\varphi = 0$ metal grounded plates



Determine $\varphi(\rho, z)$ both inside and outside the cylinder; Concentrate on $z = 0$

Warm up questions:

① What are the dimensionfull parameters?

② What are the boundary conditions?
→ what is the perpendicular directions
→ what are the parallel directions

③ What do you ^{qualitatively} expect when $L \gg R$ at $z = 0$

A worked example pg. 2

Solution: (Qualitative)

① λ , L , R , and z , ρ

at $z=0$ the solution depends on how ρ compares to L, R

② Boundary conditions:


$$\varphi = 0 \text{ at } z = \pm L/2$$

$$E_{\rho} \Big|_{R+\varepsilon} - E_{\rho} \Big|_{R-\varepsilon} = \sigma = \frac{\lambda}{2\pi R} \text{ for all } z$$

→ take z, ϕ to be parallel directions and ρ to be perp directions

③ For

$R \ll \rho \ll L$ the walls are infinitely far away, Gauss Law gives:



$$E_{\rho} = \frac{\lambda}{2\pi \rho}$$

$$\varphi = -\frac{\lambda}{2\pi} \log(\rho) + \text{Const}$$

A worked example pg. 3


Then

- As $\rho \sim L$ start to feel the effect of walls
- for $R \ll L \ll \rho$ the fields will decrease exponentially
- Inside the cylinder expect

$$E_p = 0$$

$\psi = \text{const}$ up to corrections suppressed by R^2/L^2

$$\psi(\rho) \Big|_{z=0}$$



A worked example pg. 4

Solution (Quantitative)

$$\psi = \sum_k R_k(\rho) Z_k(z)$$

The Laplace eqn inside and outside

$$-\nabla^2 \psi = 0$$

$$\left[\begin{array}{c} -\frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + \frac{\partial^2}{\partial z^2} \end{array} \right] \psi = 0$$

Leads to the separated eqs:

$$\left[\begin{array}{c} -\frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + k^2 \end{array} \right] R_k(\rho) = 0$$

$$\left[\begin{array}{c} -\frac{\partial^2}{\partial z^2} - k^2 \end{array} \right] Z_k(z) = 0$$

$$Z_k = \underbrace{A_0 + B_0 z}_{k=0 \text{ solution}} + \underbrace{\sum_k A_k \cos kz + B_k \sin kz}_{\text{general solution}}$$



Boundary conditions at $z = -L/2$ and $z = L/2$
+ symmetry $\psi(z = -L/2) = \psi(z = L/2) = 0$

$$Z_k = \sum_n A_n \cos(k_n z)$$

A worked example pg. 4

Now since the function must vanish at $z = -L/2$ and $L/2$

$$k_n = \frac{(2n+1)\pi}{L}$$



...


A worked example pg. 5

From the radial direction

$$R_k(\rho) = A_k I_0(k\rho) + B_k K_0(k\rho)$$

Asymptotics:

for $x \ll 1$:



$$I_0 = 1 + x^2/4 + \dots$$

$$K_0 = -\left[\log \frac{x}{2} + \gamma_E\right] I_0$$

+ $O(x^2)$

for $x \gg 1$

$$I_0 = \frac{e^x}{\sqrt{2\pi x}} \left(1 + O\left(\frac{1}{x}\right)\right)$$

$$K_0 = e^{-x} \sqrt{\frac{\pi}{2x}}$$

So inside the cylinder

$$R_k(\rho) = A_k I_0(k\rho)$$

And outside

$$R_k(\rho) = B_k K_0(k\rho)$$

Continuity at $\rho = R$ shows $B_k = I_0(kR)$ $A_k = K_0(kR)$

$$R_k(\rho) = C_k \left[K_0(kR) I_0(k\rho) \Theta(R - \rho) \right.$$

$$\left. + I_0(kR) K_0(k\rho) \Theta(\rho - R) \right]$$

A worked example pg. 6

So the solution at this point is

$$\varphi(\rho, z) = \sum_n C_n \left[K_0(kR) I_0(k\rho) \Theta(R-\rho) + I_0(kR) K_0(k\rho) \Theta(\rho-R) \right] \times \cos(k_n z)$$

$$\text{where } k_n = \frac{(2n+1)\pi}{L} \quad n=0, 1, 2, 3, \dots$$

From the jump condition can determine C_n

$$E_p^{\text{out}} - E_p^{\text{in}} = \frac{\lambda}{2\pi R}$$

$$E_p^{\text{in}} = -\sum_n C_n K_0'(kR) I_0'(k\rho) k_n \cos(k_n z)$$

$$E_p^{\text{out}} = -\sum_n C_n I_0(kR) K_0'(k\rho) k_n \cos(k_n z)$$

$$E_p^{\text{in}}(k) = \frac{2}{L} \int_{-L/2}^{L/2} \cos(k_n z) E_p^{\text{out}}$$

$$= -C_n K_0(kR) I_0'(kR) k_n$$

$$E_p^{\text{out}}(k) = -C_n I_0(kR) K_0'(kR) k_n$$

A worked example pg. 7

Now

$$E_p^{\text{out}}(k) - E_p^{\text{in}}(k) = \frac{2}{L} \int_{-L/2}^{L/2} \frac{\lambda}{2\pi R} \cos(k_n z) dz = \frac{\lambda \lambda}{2\pi R}$$
$$= \frac{(-1)^n 2\lambda}{(1+2n)\pi^2 R}$$

So we get that

Wronsk

$$-C_n \left[I_0(k_n R) K_0'(k_n R) - K_0(k_n R) I_0'(k_n R) \right] k_n$$
$$= \frac{(-1)^n 2\lambda}{(1+2n)\pi^2 R}$$

Recognizing the Wronskian (which often appears in jump conditions, ^{that are derived from Gauss law}) we know that this will take a simple form. From Bessel's eqn:

$$\left[-\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) + k^2 \right] (I_0(k\rho) \text{ or } K_0(k\rho)) = 0$$

$p(x)$ of Sturm-Liouville eqn $\Rightarrow p(x) = \rho$

We know that $\rho \times \text{Wronsk}(k\rho) = \text{const}$ ✓

• Use series to show $\text{Wronsk} \Big|_{\rho=R} = \frac{-1}{kR} \Big|_{\rho=R} = \frac{-1}{kR}$

A worked example pg. 8

So

$$-C_n \left[\frac{-1}{k_n R} \right] k_n = \frac{(-1)^n 2\lambda}{(1+2n)\pi^2 R}$$

So C_n is

$$C_n = \frac{(-1)^n 2\lambda}{(1+2n)\pi^2}$$

Thus we have determined the full solution

$$\psi(\rho, z) = \lambda \sum_{n=0}^{\infty} \frac{(-1)^n 2}{(1+2n)\pi^2} \times$$

$$\left[K_0(kR) I_0(k\rho) \Theta(R-\rho) + I_0(kR) K_0(k\rho) \Theta(\rho-R) \right] \\ \times \cos(k_n z)$$

Lets look at $z=0$ and outside

$$\psi(\rho) = \lambda \sum_{n=0}^{\infty} \frac{(-1)^n 2}{(1+2n)\pi^2} = I_0(kR) K_0(k\rho) \Theta(\rho-R)$$

A worked example pg. 9

• for $p > R$ but $p \ll L$ then

$$k_n p \approx \frac{(2n+1)\pi p}{L} \ll 1 \quad \text{for almost all } n$$

and $I_0 \approx 1$ $K_0 \approx -\ln k_n p - \delta_E \approx -\ln p + 2 - \delta_E + \ln k_n$

$$\varphi(p) = \lambda \sum_{n=0}^{\infty} (-1)^n \frac{2}{(1+2n)\pi^2} \left[(-\ln p) + \ln k_n + \text{const} \right]$$

$$\varphi(p) = -\frac{\lambda}{2\pi} \ln p + \text{const}$$

this was by construction

we used that $\lambda \sum_{n=0}^{\infty} (-1)^n \frac{2}{(1+2n)\pi^2} = \frac{\lambda}{2\pi}$

So

$$\varphi(p) = -\frac{\lambda}{2\pi} \ln p + \text{const} \quad \checkmark$$

• for p large $R \ll L \ll p$ find

$$K_0(k_n p) \approx \frac{1}{\sqrt{2\pi k_n p}} e^{-k_n p} \quad k_n = \frac{(2n+1)\pi p}{L}$$

• The larger the n the more it is suppressed. keep only the $n=0$ term


A worked example pg. 10

Find outside

$$\psi(\rho) = \lambda \sum_{n=0}^{\infty} \frac{(-1)^n 2}{(1+2n)\pi^2} \overbrace{I_0(kR) K_0(k\rho)}$$

$$\psi(\rho) \approx \lambda \frac{2}{\pi^2} \frac{e^{-\pi\rho/L}}{\sqrt{2(\pi\rho/L)}} \quad \left. \begin{array}{l} n=0 \\ \text{only} \\ + \text{asymptotic} \end{array} \right\}$$

$$\psi(\rho) = \lambda \frac{\sqrt{2}}{\pi^2} \frac{e^{-\pi\rho/L}}{\sqrt{\rho/L}}$$



L/R = 10

